

Q: How is PEN-Cur consumed as a Chemo/Radiation sensitizer and protector?

A: Take it 4 days before, during and 3 days after chemo/radiation treatment.

RECOMMENDED DOSAGE FOR CO-TREATMENT WITH CHEMOTHERAPY/RADIATION

30mg per kilo of your body weight - consumed once per day.

Example: If your body weight is 60 kg, therefore, $60\text{kg} \times 30\text{mg} = 1,800\text{mg}$.

Chemo

1 to 4 days before Chemo	CHEMOTHERAPY DAY 2 to 6 Hours before Chemo	1 to 3 days after chemo	4 th to 17 th day after chemo
30 minutes before meal, Best taken on an empty stomach.	Take it in the morning before breakfast or 30 minutes before meal.	30 minutes before meal, Best taken on an empty stomach.	1 to 2 teaspoons - 30 minutes before meal. (Subject to own financial ability).


Radiation

1 to 3 Days before Radiation	2 to 6 Hours before Radiation	During rest days
30 minutes before meal. Best taken on an empty stomach.	2 to 6 hours before radiation.	1 to 2 teaspoons - 30 minutes before meal. Best taken on an empty stomach.

NOTE

- A full teaspoon of PEN-Cur weighs approximately 500mg.
- **Leukemia** and **Lymphoma** cancers - take **2/3** of the recommended dose.
- Do not take **N-SAIDs** –Non steroid anti-inflammation drugs (a kind of pain killer) during **PEN-Cur treatment**, as this drug lowers the effectiveness of PEN-Cur.

Oral **PEN-Cur** Powder Preparation Instructions



Every teaspoon of PEN-Cur weighs about 500mg. Put your required dosage inside a bowl. Add in 1.5 times of drinking water carefully.



Stir slowly until it disperses into the water. Add in half a bowl of water and stir well.



Add in a little more water. Scrape the remaining PEN-Cur powder around the edges and stir into the water. It is ready to drink.